

REVTeX 4 Connection between the elastic G_{Ep}/G_{Mp} and $P \rightarrow \Delta$ form factors.

Paul Stoler*

Physics Department, Rensselaer Polytechnic Institute, Troy, NY 12180

(Dated: October 11, 2002)

It is suggested that the falloff in Q^2 of the $P \rightarrow \Delta$ magnetic form factor G_M^* is related to the recently observed falloff of the elastic electric form factor G_{Ep}/G_{Mp} . Calculation is carried out in the framework of a GPD model whose parameters are determined by fitting the elastic form factors F_{1p} and F_{2p} and isospin invariance. When applied to the $P \rightarrow \Delta$ transition with no additional parameters, the shape of G_M^* is found to exhibit the requisite falloff with Q^2 .

The $P \rightarrow \Delta(1232)$ form factor G_M^* exhibits a more rapid decrease with respect to Q^2 than is typically observed in other baryons [1] [2], such as G_{Mp} in elastic scattering from a proton, or $A_{1/2}$ in the transition $P \rightarrow S_{11}(1535)$. A recent Jefferson Lab (JLab) measurement [3] finds that the ratio G_{Ep}/G_{Mp} for elastic scattering falls with Q^2 more rapidly than previously expected. This has given rise to much theoretical activity [4] [5] to attempt to understand the underlying physics. In this note it is suggested that this behavior in G_{Ep}/G_{Mp} is related to that of G_M^* .

As a basis it is assumed that the form factor is dominated by soft mechanisms, and a GPD-handbag approach [6] [7] [8] is utilized. Form factors are the zero'th moments of the GPDs with skewedness $\xi = 0$. For elastic scattering, the Dirac and Pauli form factors are given by

$$F_1(t) = \int_{-1}^1 \sum_q H^q(x, \xi, t) dx \quad (1)$$

$$F_2(t) = \int_{-1}^1 \sum_q E^q(x, \xi, t) dx \quad (2)$$

where q signifies flavors. In the following, with $\xi=0$, for brevity the GPD's are denoted $H^q(x, t) \equiv H^q(x, 0, t)$, and $E^q(x, t) \equiv E^q(x, 0, t)$.

Resonance transition form factors access components of the GPDs which are not accessed in elastic scattering. The $N \rightarrow \Delta$ form factors are related to isovector components of the GPDs [9] [10];

$$2G_M^* = \int_{-1}^1 \sum_q H_M^q(x, t) dx \quad (3)$$

$$2G_E^* = \int_{-1}^1 \sum_q H_E^q(x, t) dx \quad (4)$$

$$2G_C^* = \int_{-1}^1 \sum_q H_C^q(x, t) dx \quad (5)$$

where G_M^* , G_E^* and G_C^* are magnetic, electric and Coulomb transition form factors [11], and H_M^q , H_E^q , and H_C^q are isovector GPDs, which can be related to elastic GPDs in the large N_C chiral limit through isospin rotations. Analogous relationships can be obtained for the $N \rightarrow S_{11}$ and other transitions. Here, the connection between GPDs involved in the elastic and $N \rightarrow \Delta$ form factors is explored to obtain the connection between the Q^2 dependence of the G_{Ep} and G_M^* .

In refs. [9] [10] it is noted that, in the large N_c limit, assuming chiral and isospin symmetry the GPDs for the $P \rightarrow \Delta(1232)$ transition are expected to be isovector components of the elastic GPDs, given by

$$H_M^{(IV)} = \frac{2}{\sqrt{3}} E_M^{(IV)} = \frac{2}{\sqrt{3}} (E_p^u - E_p^d). \quad (6)$$

E_p^u and E_p^d are the GPD's for the u and d quarks respectively. Thus the $P \rightarrow \Delta$ form factor should be obtainable by analysis of the Pauli form factor F_{2p} (eq. 2). The Dirac and Pauli form factors, F_{1p} and F_{2p} , are related to the measured Sachs form factors G_{Mp} and G_{Ep} by

$$F_{1p}(Q^2) = \frac{1}{\tau + 1} (\tau G_{Mp}(Q^2) + G_{Ep}(Q^2)) \quad (7)$$

$$F_{2p}(Q^2) = \frac{1}{\kappa(\tau + 1)} (G_{Mp}(Q^2) - G_{Ep}(Q^2)) \quad (8)$$

$$(9)$$

with $\tau = Q^2/4M_p$. To obtain E_p^u and E_p^d , needed for eq. 6, the available data for G_{Mp} and the recent JLab data [3] on G_{Ep}/G_{Mp} were fit, as reported in ref. [12], using a parameterization of the GPDs such as in [13] [14] [15] [16].

The specific functional form for $H_P(x, t)$ and $E_P(x, t)$ is a Gaussian plus small power law shape in $-t$ ($\equiv Q^2$) to account for the high the measured form factors at very high Q^2 .

$$H_P(x, t) = f(x) \exp(\bar{x}t/4x\lambda_H^2) + \dots \quad (10)$$

$$E_P(x, t) = k(x) \exp(\bar{x}t/4x\lambda_E^2) + \dots, \quad (11)$$

in which $\bar{x} \equiv 1 - x$, and $\dots$ indicates the addition of small power components added in ref. [12] to account for

*Electronic address: stoler@rpi.edu

higher Q^2 contributions to the form factors. The conditions at $Q^2=0$ are $H(x,0) = e_u f_u(x) + e_d f_d(x)$ and $E(x,0) = k_u(x) + k_d(x)$. Here, $f_u(x)$ and $f_d(x)$ are the u and d valence quark distribution functions measured in DIS. The functions $k_u(x)$ and $k_d(x)$ are not obtainable from DIS. Following ref. [15] the form used was $k^q(x) \propto \sqrt{1-x} f^q(x)$, with normalizations obtained employing isospin symmetry, and by requiring the proton and neutron form factors to have their known values near $Q^2=0$, that is $F_{1p}(0) = 1$, $F_{2p}(0) = 1.79$, $F_{1n}(0) = 0$, $F_{2n}(0) = -1.91$. This gives

$$F_{2u}(0) \equiv \kappa_u = \int k_u(x) dx = 1.67 \quad (12)$$

$$F_{2d}(0) \equiv \kappa_d = \int k_d(x) dx = -2.03 \quad (13)$$

and

$$F_{1u}(0) = \int e_u f_u(x) dx = 2/3 \quad (14)$$

$$F_{1d}(0) = \int e_d f_d(x) dx = 1/3 \quad (15)$$

Adequate fits to G_{Mp} and G_{Ep}/G_{Mp} , or equivalently F_{1p} and F_{2p}/F_{1p} , were obtained with $\lambda_H = 0.76$ GeV/c and $\lambda_E = 0.67$ GeV/c. The results are shown in figs. 1 and 2.




FIG. 1: Dirac form factor $F_{1p}(Q^2)$ relative to the dipole $G_D = 1/(1 + Q^2/.71)^2$. The data are extracted using the recent JLab. data [3] for G_{Ep}/G_{Mp} , and a recent reevaluation [20] of SLAC data of G_{Mp} [18] [19]. The curve is the result of the fit as discussed in the text.

The resulting E_p^u and E_p^d were inserted into eq. 6 to obtain an estimate for G_M^* . At $Q^2=0$ one gets $G_M^*(0) = 2.14$, which is somewhat lower than the experimental value of $G_M^*(0) \sim 3$. Such a disagreement is not surprising [9] [10] given the very approximate nature of eq. 6. The obtained G_M^* was overall renormalized to take this ratio into account, and the result is shown in fig. 3.




FIG. 2: The Pauli form factor $F_2/1.79F_D$ relative to the dipole $F_D = 1/(1 + Q^2/.71)^2$. The data are extracted using the recent JLab. data [3] for F_{2p}/F_{1p} , multiplied by the fit curve for F_{1p}/F_D shown in fig. 1. The curve is the result of the simultaneous fit to the G_{Ep}/G_{Mp} and G_{Mp} data as discussed in the text and fig. 1.




FIG. 3: The $N \rightarrow \Delta$ magnetic form factor $G_M^*(Q^2)/3G_D$ relative to the dipole $G_D = 1/(1 + Q^2/.71)^2$. The data are a compendium of world data by ref [21]. The curve is the result of the procedures discussed in the text.

The similar shapes of the curves in figs. 2 and 3 can be ascribed to their connection via eq. 6. This can be understood by the observation that F_2 is nearly all isovector spin-flip, as is the G_M^* . However, the difference in the mass of the $\Delta(1232)$ and the nucleon, which is a measure of the SU3 symmetry breaking, and the fact that F_1 also has an isovector component would make the observed non-negligible differences in the normalization not surprising.

Although this note suggests a common physical origin in the the Q^2 behavior of G_{Ep}/G_{Mp} and G_M^* , a complete understanding will require theoretical treatments based on rigorous and consistent relativistic treatment which are beyond the scope of this communication.

Acknowledgments: The author thanks G.A. Miller,

A.V. Radyushkin and M. Vanderhaeghen for helpful discussions. The work was partially supported by the *National Science Foundation*.

-
- [1] P. Stoler, *Physics Reports* **226**, 103 (1993).
 - [2] V.V. Frolov *et al.*, *Phys.Rev.Lett.* **82**,45 (1999).
 - [3] M.K. Jones *et al.* *Phys. Rev. Lett.* **84**,1398 (2000); O. Gayou *et al.* *Phys. Rev.* **C64**,038202 (2001).
 - [4] G.A. Miller and M.R. Frank, *Phys. Rev.*,**C65**, 065205 (2002). nucl-th/0206027; G.A. Miller,*Phys. Rev.*,**C66**, 03220(R) (2002). nucl-th/0207007;
 - [5] J.P. Ralston and P. Jain, Proceedings of the Workshop on Exclusive Processes at High Momentum Transfer, A. Radyushkin and P. Stoler, eds., World Scientific, Singapore, 105; VII International Conference on the Intersections of Particle and Nuclear Physics (Quebec City 2000). Z. Parsehand W. Marciano, eds., American Institute of Physics, (2000).
 - [6] X. Ji, *Phys. Rev. Lett.* **78**, 610 (1997).
 - [7] A.V. Radyushkin, *Phys. Lett.* **B380**,417 (1996); *Phys. Rev.* **D56**,5524 (1997).
 - [8] J. Collins, L. Frankfurt, and M. Strikman, *Phys. Rev.*, **D56**, 2982 (1997).
 - [9] L.L.Frankfurt *et al.*, *Phys. Rev. Lett.* **84**,2589 (2000).
 - [10] K.Goeke, M.V. Polyakov, and M. Vanderhaeghen, *Prog. Part. Nucl. Phys.* **47**, 401,2001 (2001).
 - [11] H.F. Jones and M.D. Scadron, *Annals of Physics* **81**, 1 (1979).
 - [12] P. Stoler, *Phys. Rev.* **D65**, 053013 (2002).
 - [13] A.V. Radyushkin, *Phys. Rev.* **D58**,114008 (1998).
 - [14] M. Diehl, Th. Feldmann, R. Jakob and P. Kroll, *Eur. Phys.* **C8**, 409 (1999); M. Diehl, Th. Feldmann, R. Jakob and P. Kroll, *Nucl. Phys.* **B596**, 33 (2001), Erratum-ibid. **B605**, 647 (2001).
 - [15] A. Afanasev, E-print: hep-ph/9910565;
 - [16] M. Burkardt, Proceedings of the Workshop on Exclusive Processes at High Momentum Transfer, A. Radyushkin and P. Stoler, eds., World Scientific, Singapore, 99 (2003), and references within.
 - [20] E.J. Brash *et al.*, *Phys. Rev.* **C65**, 051001(R) (2002).
 - [18] R.G. Arnold *et al.*, *Phys. Rev. Lett.* **57**, 174 (1986).
 - [19] L. Andivahis *et al.*, *Phys. Rev.* **D50**, 5491 (1994).
 - [20] E.J. Brash *et al.*, *Phys. Rev.* **C65**, 051001(R) (2002).
 - [21] S.S. Kamalov and S.N. Yang, *Phys. Rev. Lett.* **83**, 4494 (1999).